

Knee Junction Dilemma

Out of the box solutions for challenging knees

BIBLIOGRAPHY

1. Eckenrode, Brian J., Kietrys David M., Parrott, J. Scott. Effectiveness of Manual Therapy on Pain and Self-Reported Function in Individuals with Patellofemoral Pain: Systematic Review and Meta-Analysis. *J Orthop Sports Phys Ther*:1-48. Epub 6 Jan 2018. <https://doi.org/10.2519/jospt.2018.7243> 6 Jan 2018.
2. Sebro R, Weintraub S. Association between lateral patellar osteoarthritis and knee morphology and alignment in young adults. *Clinical Radiology*. 2017;72(9). doi: 10.1016/j.crad.2017.03.008.
3. Collins CK, Johnson R, Masaracchio M. A Clinical Reasoning Model for Manual Physical Therapy. *Orthopedic Physical Therapy Practice*. 2017;29(4).
4. Sihvonen R, Paavola M, Malmivaara A, et al. Arthroscopic partial meniscectomy versus placebo surgery for a degenerative meniscus tear: a 2-year follow-up of the randomised controlled trial. *Annals of the Rheumatic Diseases*. 2017. doi:10.1136/annrheumdis-2017-211172.
5. Takacs J, Krowchuk NM, Garland SJ, Carpenter MG, Hunt MA. Dynamic Balance Training Improves Physical Function in Individuals with Knee Osteoarthritis: A Pilot Randomized Controlled Trial. *Archives of Physical Medicine and Rehabilitation*. 2017;98(8):1586-1593. doi: 10.1016/j.apmr.2017.01.029.
6. Lai CC¹, Ardern CL^{2,3,4}, Feller JA⁵, Webster KE². Eighty-three per cent of elite athletes return to preinjury sport after anterior cruciate ligament reconstruction: a systematic review with meta-analysis of return to sport rates, graft rupture rates and performance outcomes. *Br J Sports Med*. 2017 Feb 21. pii: bjsports-2016-096836. doi: 10.1136/bjsports-2016-096836.
7. Nascimento LR^{1,2}, Teixeira-Salmela LF¹, Souza RB¹, Resende RA¹.. Hip and Knee Strengthening is More Effective Than Knee Strengthening Alone for Reducing Pain and Improving Activity in Individuals with Patellofemoral Pain: A Systematic Review with Meta-Analysis. *J Orthop Sports Phys Ther*. 2017 Oct 15:1-35. doi: 10.2519/jospt.2018.7365
8. Hamstra-Wright KL¹, Earl-Boehm J, Bolgia L, Emery C, Ferber R.. Individuals with Patellofemoral Pain Have Less Hip Flexibility Than Controls Regardless of Treatment Outcome. *Clin J Sport Med*. 2017 Mar;27(2):97-103. doi: 10.1097/JSM.0000000000000307.
9. Knee Ligament Sprain Guidelines: Revision 2017: Using the Evidence to Guide Physical Therapist Practice Published: *Journal of Orthopaedic & Sports Physical Therapy*, 2017 Volume:47 Issue:11 Pages:822–823 DOI:10.2519/jospt.2017.0510
10. Kwak YH¹, Lee S¹, Lee MC¹, Han HS². Large meniscus extrusion ratio is a poor prognostic factor of conservative treatment for medial meniscus posterior root tear. *Knee Surg Sports Traumatol Arthrosc*. 2017 Feb 15. doi: 10.1007/s00167-017-4441-x.

11. Paterson K, Kasza J, Hunter D, et al. Longitudinal association between foot and ankle symptoms and worsening of symptomatic radiographic knee osteoarthritis: data from the osteoarthritis initiative. *Osteoarthritis and Cartilage*. 2017;25(9):1407-1413. doi: 10.1016/j.joca.2017.05.002.
12. Ahn JH1, Jeong HJ, Lee YS, Park JH, Lee JW, Park JH, Ko TS. Comparison between conservative treatment and arthroscopic pull-out repair of the medial meniscus root tear and analysis of prognostic factors for the determination of repair indication. *Arch Orthop Trauma Surg*. 2015 Jul 5.
13. Plinsinga ML1, van Wilgen CP2,3,4, Brink MS5, Vuvan V1, Stephenson A1, Heales LJ1,6, Mellor R1, Coombes BK7, Vicenzino BT1. Patellar and Achilles tendinopathies are predominantly peripheral pain states: a blinded case control study of somatosensory and psychological profiles. *Br J Sports Med*. 2017 Jul 11. pii: bjsports-2016-097163. doi: 10.1136/bjsports-2016-097163.
14. Elmansori A1,2, Lording T3, Dumas R2, Elmajri K1,2, Neyret P1,2, Lustig S4,5. Proximal tibial bony and meniscal slopes are higher in ACL injured subjects than controls: a comparative MRI study. *Knee Surg Sports Traumatol Arthrosc*. 2017 Feb 17. doi: 10.1007/s00167-017-4447-4
15. Steidle-Kloc E1, Culvenor AG1,2, Dörrenberg J1, Wirth W1, Ruhdorfer A1, Eckstein F1. Relationship between knee pain and infra-patellar fat pad morphology - A within- and between-person analysis from the Osteoarthritis Initiative. *Arthritis Care Res (Hoboken)*. 2017 Jul 13. doi: 10.1002/acr.23326.
16. Wikstrom EA1, Song K2, Pietrosimone BG2, Blackburn JT2, Padua DA2. Visual Utilization During Postural Control in Anterior Cruciate Ligament Deficient & Reconstructed Patients: Systematic Reviews & Meta-Analyses. *Arch Phys Med Rehabil*. 2017 May 5. pii: S0003-9993(17)30302-7. doi: 10.1016/j.apmr.2017.04.010.
17. Authors: Allison R. Toole, PT, DPT, OCS1, Matthew P. Ithurburn, PT, PhD, DPT, OCS2, Mitchell J. Rauh, PT, PhD, MPH, FACSM3, Timothy E. Hewett, PhD, FACSM4, Mark V. Paterno, PT, PhD, SCS5–7, Laura C. Schmitt, PT, PhD8–10 Young Athletes Cleared for Sports Participation After Anterior Cruciate Ligament Reconstruction: How Many Actually Meet Recommended Return-to-Sport Criterion Cutoffs? Published: *Journal of Orthopaedic & Sports Physical Therapy*, 2017 Volume:47 Issue:11 Pages:825–833 DOI:10.2519/jospt.2017.7227
18. Int J Rheum Dis. 2015 Jul 14. doi: 10.1111/1756-185X.12696. Does knee malalignment predict the efficacy of realignment therapy for patients with knee osteoarthritis? Rezaeian ZS1,2,3, Smith MM4, Skaife TL5, Harvey WF6, Gross KD7, Hunter DJ2,3.
19. Daniel Ramskov, PT, MHSc1, Christian Barton, PT, PhD3–6, Rasmus O. Nielsen, PT, MHSc1,2, Sten Rasmussen, MD2,7,8 High Eccentric Hip Abduction Strength Reduces the Risk of Developing Patellofemoral Pain Among Novice Runners Initiating a Self-Structured Running Program: A 1-Year Observational Study Published: *Journal of Orthopaedic & Sports Physical Therapy*, 2015, Volume: 45 Issue: 3 Pages: 153-61 doi:10.2519/jospt.2015.5091
20. Anterior Cruciate Ligament Injury in Male and Female Athletes: A Prospective Study. . *Am J Sports Med*. 2015 Dec

21. Mouton C1, Seil R1,2, Meyer T3, Agostinis H1, Theisen D4 Combined anterior and rotational laxity measurements allow characterizing personal knee laxity profiles in healthy individuals. *Knee Surg Sports Traumatol Arthrosc.* 2015 Dec;23(12):3571-7. doi: 10.1007/s00167-014-3244-6. Epub 2014 Aug 26.
22. Hagino T1, Ochiai S, Senga S, Yamashita T, Wako M, Ando T, Haro H. Meniscal tears associated with anterior cruciate ligament injury. *Arch Orthop Trauma Surg.* 2015 Aug 19.
23. Grooms D1, Appelbaum G, Onate J. Neuroplasticity following anterior cruciate ligament injury: a framework for visual-motor training approaches in rehabilitation. *J Orthop Sports Phys Ther.* 2015 May;45(5):381-93. doi: 10.2519/jospt.2015.5549. Epub 2015 Jan 10.
24. Shimamura K. et al. Regional Interdependence of the Hip and Lumbopelvic Region in Division II Collegiate Level Baseball Pitchers: A Preliminary Study. *Int Jnl Sports Phys Ther.* 2015;10(1):1-12.
25. Park do Y1, Min BH2, Choi BH3, Kim YJ4, Kim M4, Suh-Kim H5, Kim JH6 The Degeneration of Meniscus Roots Is Accompanied by Fibrocartilage Formation, Which May Precede Meniscus Root Tears in Osteoarthritic Knees. *Am J Sports Med.* 2015 Dec;43(12):3034-44. doi: 10.1177/0363546515605086. Epub 2015 Oct.
26. Umegaki H, Ikezoe T, Nakamura M, et al. The effect of hip rotation on shear elastic modulus of the medial and lateral hamstrings during stretching. *Manual Therapy.* 2015;20(1):134-137. doi: 10.1016/j.math.2014.07.016.
27. Hirschmann A1,2, Buck FM3, Herschel R4, Pfirrmann CW3, Fucentese SF4. Upright weight-bearing CT of the knee during flexion: changes of the patellofemoral and tibiofemoral articulations between 0° and 120°. *Knee Surg Sports Traumatol Arthrosc.* 2015 Nov 4.
28. Khayambashi, Khalil, et al. "Hip Muscle Strength Predicts Noncontact Anterior Cruciate Ligament Injury in Male and Female Athletes." *The American Journal of Sports Medicine*, vol. 44, no. 2, 8 Dec. 2015, pp. 355–361, <https://doi.org/10.1177/0363546515616237>.
29. *Arch Phys Med Rehabil.* 2014 Feb 14. pii: S0003-9993(14)00116-6. doi: 10.1016/j.apmr.2014.01.025. Comparative evaluation of core muscles recruitment pattern in response to sudden external perturbations in patients with patellofemoral pain syndrome and healthy subjects. Shirazi ZR1, Moghaddam MB2, Motealleh A3.
30. Berger MJ. The relationship between Quadriceps muscle volume and patellar cartilage morphology in patients with knee osteoarthritis. *Journal of Exercise, Sports & Orthopedics.* 2014;1(3). doi:10.15226/2374-6904/1/3/00116.
31. Sueki D, Cleland J, Wainner R. A Regional Interdependence Model of Musculoskeletal Dysfunction: Research, Mechanisms, and Clinical Implications. *Jnl Manual and Manipulative Ther.* 2013;21(2):90-103.
32. Brummitt, J. The Role of Massage in Sports Performance and Rehabilitation: Current Evidence and Future Direction. *N Am J Sports Phys Ther.* 2008 Feb; 3(1): 7–21.

33. Wainer R. et al. Regional Interdependence: A Musculoskeletal Model Whose Time has Come. *Journal of Orthopaedic & Sports Physical Therapy*, 2007; 37(11): 658-660
34. Chmielewski TL, Hodges MJ, Horodyski M, et al. Investigation of clinical agreement in evaluating movement quality during unilateral lower extremity functional tasks: a comparison of 2 rating methods. *JOSPT*. 2007; 37:1222-9.
35. Loudon JK, D Wiesner, HL Goist-Foley, C Asjes, KL Loudon. Intrarater reliability of functional performance tests for subjects with patellofemoral pain syndrome. *J Athl Training*. 2002; 37:256-61